



doi:10.1016/j.gca.2004.03.005

Lipid biomarkers and carbon-isotopes of modern travertine deposits (Yellowstone National Park, USA): Implications for biogeochemical dynamics in hot-spring systems

CHUANLUN L. ZHANG,^{1,*} BRUCE W. FOUKE,² GEORGE T. BONHEYO,² AARON D. PEACOCK,³ DAVID C. WHITE,³ YONGSONG HUANG,⁴ and CHRISTOPHER S. ROMANEK¹

¹Savannah River Ecology Laboratory, University of Georgia, Drawer E, Aiken, SC 29802, USA

²Department of Geology, University of Illinois, 245 Natural History Building, Urbana, IL 61801, USA

³Center for Biomarker Analysis, The University of Tennessee, Knoxville, TN 37932, USA

⁴Department of Geological Sciences, Brown University, Providence, RI 02912, USA

Abstract—Lipid biomarkers and ^{13}C fractionation patterns were used to understand the dynamics of carbon cycling during microbial metabolisms in different environments of travertine precipitation (called facies) at Spring AT-1 on Angel Terrace in the Mammoth Hot Springs complex of Yellowstone National Park, USA. Microbial mats that encrust travertine deposits were collected for analyses of lipid biomarkers and carbon isotopes along the continuous drainage outflow system of Spring AT-1. The spring water exhibits a continuous temperature drop from 71°C in the vent at top to 24°C in the distal slope at bottom. Phospholipid fatty acids (PLFA) and glycolipid fatty acids (GLFA) exhibit distinctly different compositions in each of the facies, which are consistent with partitioning of the bacterial 16S rRNA gene sequences in the Spring AT-1 travertine facies (Fouke et al., 2003).

The $\delta^{13}\text{C}$ composition of total biomass within the microbial mats decreases from -16.1‰ in the vent to -23.5‰ in the distal slope. However, lower values occur in the pond (-26.0‰) and the proximal slope (-28.0‰) between the vent and the distal slope. Isotopic compositions of PLFA and GLFA have variations similar to those of total biomass. The average $\delta^{13}\text{C}$ values of PLFA are $-12.4 \pm 5.2\text{‰}$ ($n = 10$ individual fatty acids, same below) in the vent, $-33.0 \pm 3.1\text{‰}$ ($n = 11$) in the pond, $-33.7 \pm 3.8\text{‰}$ ($n = 16$) in the proximal slope, and $-22.4 \pm 3.4\text{‰}$ ($n = 10$) in the distal slope; the average $\delta^{13}\text{C}$ values of GLFA are $-19.6 \pm 3.0\text{‰}$ ($n = 3$) in the vent, $-30.4 \pm 4.7\text{‰}$ ($n = 8$) in the pond, $-36.9 \pm 2.8\text{‰}$ ($n = 12$) in the proximal slope, and $-27.9 \pm 3.1\text{‰}$ ($n = 13$) in the distal slope. In particular, fatty acids in the vent are enriched in ^{13}C relative to the total biomass, which is consistent with the notion that the biosynthetic pathways of the extant microbial community in the vent may be dominated by *Aquificales* using the reversed tricarboxylic acid cycle. Fractionations between fatty acids and total biomass in the pond, the proximal slope and the distal slope suggest the involvement of other biosynthetic pathways for CO_2 fixation by extant microbial populations. The results indicate that lipid biomarkers provide valuable information on the changing diversity and activity of microbial communities in different depositional environments. Carbon-isotope fractionations, on the other hand, can provide insight into the operating biosynthetic pathways associated with different organisms in the changing environment. This integrated approach may serve as a powerful tool for identifying functional metabolism within a community and identify shifts in microbial community structure in modern hot-spring systems. Copyright © 2004 Elsevier Ltd

Keywords: Phospholipid fatty acids, glycolipid fatty acids, lipid biomarkers, stable carbon isotopes, *Aquificales*, cyanobacteria, green sulfur bacteria, green non-sulfur bacteria, Angel Terrace, Mammoth Hot Springs, Yellowstone National Park

INTRODUCTION

Lipid biomarkers provide quantitative information about the structure of extant microbial communities without the need for culturing and isolation (White, 1988). Lipids are also one of the most useful biochemical measures of in situ interactions between microbial species and their environments because lipid compositions can indicate temperature-, redox-, stress-, or nutritional conditions (Ray et al., 1971; Fork et al., 1979; Ringelberg et al., 1989; Jahnke, 1992). Furthermore, bacteria commonly have ester-linked lipids and some bacterial species contain additional non-phytanyl ether lipids (Langworthy et al., 1983; De Rosa et al., 1988; Huber et al., 1992). Archaea, on the other hand, have phytanyl ether lipids (Langworthy et al., 1983). Thus, analysis of lipid structures permits identification

of these two microbial domains in unknown environmental samples.

Different microorganisms have different biosynthetic pathways. In particular, autotrophic microorganisms can use several pathways for CO_2 fixation (Ratledge and Wilkinson, 1988). A variety of phototrophs, which include cyanobacteria, use the Calvin cycle, which requires the key enzymes of ribulose 1,5-bisphosphate carboxylase/oxygenase (RubisCO) and ribulokinase. Methanogens and several groups of known chemolithoautotrophic bacteria use the acetyl-CoA pathway for CO_2 fixation (Fuchs, 1989; Vorholt et al., 1995). Other bacteria, such as *Hydrogenobacter* and *Aquifex* are known to use the reversed tricarboxylic acid (rTCA) cycle for CO_2 fixation (Fuchs, 1989; Beh et al., 1993). Another pathway is the 3-hydroxypropionate (HP) cycle, which fixes HCO_3^- rather than CO_2 . This pathway has been observed in green non-sulfur bacteria (Strauss and Fuchs, 1993) and Archaea (Menendez et al., 1999). It is suggested that the latter three pathways require

* Author to whom correspondence should be addressed (zhang@srel.edu).

less ATP for biosynthesis than the Calvin cycle (Karl, 1995); thus they may have significant ecological and evolutionary implications for life that survives under extreme conditions.

Different biosynthetic pathways may result in distinct isotope-fractionation patterns. Microorganisms using the Calvin cycle normally produce fractionations about -26‰ between biomass and CO_2 (O'Leary, 1988; Sirevåg et al., 1977; Madigan et al., 1989; Sakata et al., 1997; Popp et al., 1998); those using the acetyl-CoA pathway can produce fractionations as low as -36‰ (Preuß et al., 1989). On the other hand, the rTCA cycle and the 3-HP pathway can produce fractionations greater than (more positive) -17‰ between biomass and CO_2 (Quandt et al., 1977; van der Meer et al., 2001; Zhang et al., 2002; Londry et al., 2004).

Distinct fractionations have also been observed between lipid biomarkers and biomass for CO_2 fixation pathways used by different organisms. For example, fractionations associated with the Calvin cycle range from -7.6 to -9.9‰ between fatty acids and biomass for a cyanobacterium (Sakata et al., 1997). Fractionations associated with the 3-HP pathway range from 0.2 to -1.9‰ for a green non-sulfur bacterium (van der Meer et al., 2001). Fractionations associated with rTCA cycle in *Aquificales* and green sulfur bacteria range from 2.0 – 16.0‰ with fatty acids being enriched in ^{13}C relative to biomass (van der Meer et al., 1998; Jahnke et al., 2001; Zhang et al., 2002); however, a fractionation value of -11.8‰ has recently been reported for a sulfate-reducing bacterium (*Desulfobacter hydrogenophilus*) using rTCA for CO_2 fixation (Londry et al., 2004). These results illuminate the potential for using lipid biomarkers and molecular stable isotopes to understand diverse CO_2 fixation pathways and their effects on carbon flow associated with or mediated by microorganisms.

Biomarkers have been widely utilized as indicators of past biologic activities on Earth (Mackenzie et al., 1982; Simoneit, 1986; Ward et al., 1989; Brassell, 1992; Summons et al., 1996; Freeman, 2001; Hayes, 2001; Simoneit, 2002). The study of lipid biomarkers in hot-spring systems has particular implications for the evolution of life because the earliest common ancestor of life on earth might have evolved from a thermophilic organism (Woese, 1987; Barns and Nierzwicki-Bauer, 1997).

The present study focused on Spring AT-1 on Angel Terrace in the Mammoth Hot Springs complex of Yellowstone National Park, USA. The depositional facies and associated water chemistry, mineralogy and phylogenetic diversity have been characterized (Fouke et al., 2000, 2003). Our goal in this study was to delineate the carbon cycling pathways in each of the five different travertine depositional facies that occur along the outflow of Spring AT-1, and to use Spring AT-1 as a model for understanding microbial-biogeochemical dynamics in modern and perhaps ancient hot-spring travertine deposits.

Site Description and Previous Studies of Chemistry and Microbiology

Angel Terrace is located in the New Highland Terrace area within the Mammoth Hot Spring complex (Allen and Day, 1935; Bargar, 1978; Fouke et al., 2000). The hydrologic system at Angel Terrace is dynamic with multiple vents appearing, sealing, and reopening at a frequency of months to tens of years (Friedman, 1970; Bargar, 1978; Sorey and Colvard, 1997).





Figure 1. (Upper) Photograph of Angel Terrace Spring AT-1 (taken on July 7, 2001). (Lower) A cross section indicating facies distribution (Table 2) and flow directions (from Fouke et al., 2000).

Fouke et al. (2000) characterize Angel Terrace deposits into multiple facies based on water temperature, chemistry, mineralogy, physical structure, and travertine accumulation rates. These facies include the vent (71 – 73°C), the apron and channel (69 – 74°C), the pond (30 – 71°C), the proximal slope (28 – 54°C), and the distal slope (28 – 30°C) (Fig. 1). As the spring water cools, CO_2 degasses and the water chemistry changes from pH ~ 6 in the vent to pH 8 or higher in the distal slope (Fouke et al., 2000). As a result, total dissolved inorganic carbon (DIC) decreases from ~ 190 mg/L in the vent to ~ 70 mg/L in the distal slope. On the other hand, the carbon-isotope composition of DIC increases from $\sim -1\text{‰}$ in the vent to $\sim 5\text{‰}$ in the distal slope (Fouke et al., 2000).

Changes in the microbial communities inhabiting the outflow drainage systems of Angel Terrace are first described by Farmer and Des Marais (1994) and substantiated by Fouke et al. (2003). In particular, Fouke et al. (2003) observe that the phylogenetic diversity of bacteria is strongly partitioned among five different travertine depositional facies in the surface drainage system of the Angel Terrace (Table 1). For example, the vent is dominated by bacterial sequences affiliated with *Aquificales*; whereas, other facies are dominated by sequences affiliated with bacteria including Bacteroides, Cytophagales, Flexibacter, Firmicutes, OP11, OP8, Planctomycetales, Proteobacteria, Thermus-Deinococcus, and Verrucomicrobia (Table 1). The relative proportions of gene sequences for photo-

Table 1. Percentage of gene sequences for chemolithoautotrophs (*Aquificales*), photoautotrophs (green-sulfur bacteria, green non-sulfur bacteria, and cyanobacteria), and other bacterial species recovered from surface environments (facies) at Angel Terrace travertine in Yellowstone (Fouke et al., 2003).

Division	Facies				
	Vent	Apron and channel	Pond	Proximal slope	Distal slope
Aquificales	91	25	7	15	5
Green nonsulfur bacteria	3	4	1	16	ND
Green sulfur bacteria	<1	4	1	9	3
Cyanobacteria	1	ND	9	2	18
Other bacteria ^a	3	67	72	42	74

ND = not detected.

^a Other bacteria include BCF (Bacteroides, Cytophagales, and Flexibacter), Firmicutes, OP11, OP8, Planctomycetales, Proteobacteria, Thermus-Deinococcus, and Verrucomicrobia in different proportions in each facies, which are either heterotrophs or undetermined.

autotrophs such as green sulfur bacteria, green non-sulfur bacteria, and cyanobacteria vary from facies to facies (Table 1).

Fouke et al. (2003) also observe that there is little down stream transport of bacterial cells at the Angel Terrace Spring AT-1. This indicates that the distribution of microorganisms may be directly controlled by the physical and chemical conditions of each depositional facies. Results of this study support this hypothesis and indicate that Spring AT-1 would be an excellent natural laboratory for studying biogeochemical dynamics in the hot spring environments. Specifically, lipid biomarkers and their molecular isotope compositions would allow us to delineate pathways for CO₂ fixation within a mat community and the shifts in microbial community structure from facies to facies.

MATERIAL AND METHODS

3.1. Sample Collection

Microbial mats or fresh travertine accumulations (carbonate rocks in the vent and the apron and channel) were collected at Angel Terrace

Spring AT-1 during daylight-hours in July 2001. The samples were collected from the same spring reported by Fouke et al. (2003) but not from the exact locations. This is because the flow of the spring water is dynamic and changes pathways frequently, which results in the formation of the same facies in different locations and times. The temperature in the same facies, however, is comparable between locations, which makes this study relevant to that of Fouke et al. (2003).

In each facies, the mat material was physically peeled off and collected into 50-mL sterile plastic tubes using forceps. Non-calcified mats grew very thin in the vent and in the apron and channel because of high rates of travertine precipitation. As a result, a piece of the fresh carbonate rock was cut using a knife and transferred into the plastic tube using forceps. Two samples (A and B) were collected from each of the five selected facies (Table 2). Water temperature and pH were measured at the time of collection using a combination probe. Samples were stored on ice in the field (<10 h) and frozen immediately on dry ice at the end of the day. Samples were transported to the home institution on dry ice and lyophilized before lipid extraction and carbon-isotope determinations.

3.2. Lipid Extraction and Fatty Acid Identification

Freeze-dried materials were extracted following White et al. (1979) and (Zhang et al., 2002, 2003). The procedure employed a single-phase organic solvent system comprised of chloroform, methanol, and aqueous 50-mM phosphate buffer (pH 7.4) in the ratio of 1:2:0.8 (v:v:v). In this study, samples were powdered and a portion of the material (0.34–1 g) was used for lipid extraction. After extraction overnight, chloroform and nano-pure water were added to the extractant in equal volumes, which resulted in a two-phase system. The lower phase contained the lipids, which were collected and fractionated on a silicic acid column into neutral lipids, polar lipids (phospholipids), and glycolipids (Guckert et al., 1985). The phospholipids and glycolipids were treated using a mild alkaline methanolysis to produce fatty acid methyl esters (FAMES), namely the phospholipids fatty acids (PLFA) and glycolipid fatty acids (GLFA).

An Agilent 6890 series gas-chromatograph (GC) interfaced to an Agilent 5973 mass selective detector was used for the identification of FAMES. The GC used a 60-m non-polar column (0.25-mm I.D., 0.25- μ m film thickness) and the injector and detector were maintained at 230°C and 300°C, respectively. The column-temperature was programmed from 60°C for 1 min, ramping at 20°C/min to 150°C and holding for 4 min. This was followed by ramping at 7°C/min to 230°C and holding for 2 min, and finally ramping at 10°C/min to 300°C and holding for 3 min. Mass-spectra were determined by electron impact at 70 eV. Methyl Heneicosanoate was used as the internal standard. PLFA and GLFA were expressed as equivalent peaks against the internal standard. Double-bond positions of monounsaturated FAME were determined by GC-mass-spectrometry (MS) analysis of the dimethyl

Table 2. Water chemistry and mineralogy of Angel Terrace Spring AT-1.

Parameter	Facies				
	Vent	Apron & Channel	Pond-2	Proximal Slope-2	Distal Slope
Temp. (°C) ^a	71.4 ± 0.5	66.8 ± 0.3	50.9 ± 0.6	46.8 ± 0.5	24.3 ± 0.0
pH ^a	6.3 ± 0.1	6.7 ± 0.1	7.5 ± 0.0	8.1 ± 0.0	8.4 ± 0.0
DIC (mg/L) ^b	192	111	135	104	75
$\delta^{13}\text{C}_{\text{DIC}}$ (‰) ^b	-0.9	1.0	0.4	1.6	4.6
$\delta^{13}\text{C}_{\text{CO}_2\text{-d}}$ (‰) ^c	-5.4	-3.9	-5.9	-5.1	-4.5
$\delta^{13}\text{C}_{\text{TB}}$ (‰)	-16.1 ± 4.2	ND ^d	-26.3	-28.0 ± 0.6	-23.5 ± 0.0
Flow rate ^b	High	High	Low	Moderate	Low
Mineralogy ^b	Aragonite	Aragonite	Aragonite/Calcite	Aragonite/Calcite	Calcite
Mat color	Tan-gray travertine	Pink-white	White/green	Dark green/orange-brown	Orange

^a Mean ± 1 standard deviation for two or three replicate measurements.

^b Data are from Fouke et al. (2000). DIC = dissolved inorganic carbon.

^c Isotopic compositions of dissolved CO₂ ($\delta^{13}\text{C}_{\text{CO}_2\text{-d}}$) was calculated from $\delta^{13}\text{C}_{\text{HCO}_3}$ ($\equiv \delta^{13}\text{C}_{\text{DIC}}$) using the equations of $\delta^{13}\text{C}_{\text{CO}_2\text{-d}} = \varepsilon_{(\text{CO}_2\text{-HCO}_3)} \times (1000 + \delta^{13}\text{C}_{\text{HCO}_3})/1000 + \delta^{13}\text{C}_{\text{HCO}_3}$ and $\varepsilon_{(\text{CO}_2\text{-HCO}_3)} = 24.12 - 9866/T$, where ε is the fractionation between dissolved CO₂ and HCO₃⁻ and T is absolute temperature (Mook et al., 1974).

^d Not determined.




Figure 2. Gas chromatograph of phospholipid fatty acids from a vent sample collected from the Angel Terrace hot-spring complex in Yellowstone National Park (see Fig. 2 for description). Peak identifications are: 1, i15:0; 2, a15:0; 3, i16:0; 4, 16:0; 5, i17:0; 6, a17:0; 7, 17:0; 8, i18:0; 9, 18:1 ω 9c; 10, 18:1 ω 7c; 11, 18:1 ω 7t; 12, 18:0; 13, cy19:0; 14, 20:1 ω 9c; 15, 20:1 ω 9t; 16, 20:0; 17, 21:0 (internal standard).

disulfide adducts (Nichols et al., 1986). *Cis* and *trans* isomers of compounds were identified by known standards. Figure 2 shows an example of fatty-acid profiles for the vent facies.

3.3. Stable Carbon Isotopes

Carbon-isotope compositions of the FAMES for PLFA and GLFA were determined following Zhang et al. (2002) using a HP 6890 gas-chromatograph connected to a Finnigan MAT Delta⁺-XL mass-spectrometer. Measurements were corrected for the methyl moiety according to Abrajano et al. (1994) using the following equation:

$$\delta^{13}\text{C}_{\text{FA}} = [(\text{Cn} + 1) \times \delta^{13}\text{C}_{\text{FAME}} - \delta^{13}\text{C}_{\text{MeOH}}] / \text{Cn}$$

where $\delta^{13}\text{C}_{\text{FA}}$ is the $\delta^{13}\text{C}$ of the fatty acid, Cn is the number of carbons in the fatty acid, $\delta^{13}\text{C}_{\text{FAME}}$ is the $\delta^{13}\text{C}$ of the methylated fatty acid, and $\delta^{13}\text{C}_{\text{MeOH}}$ is the $\delta^{13}\text{C}$ of the methanol used for the methylation reaction.

Carbon-isotope compositions of total biomass were determined using bulk samples after dissolution of the associated carbonate in 1% EDTA. The residual material was then rinsed with distilled water and dried in an oven at 50°C overnight. The $^{13}\text{C}/^{12}\text{C}$ ratio of total biomass was then determined on a Delta Plus isotope ratio mass spectrometer interfaced with an Elemental Analyzer.

RESULTS

4.1. General Chemistry

Water temperature decreased from 71.4°C in the vent to 24.3°C in the distal slope at the time of sampling. The pH increased from 6.3 in the vent to 8.4 in the distal slope (Table 2). These measurements are in the same range as previous recordings (Fouke et al., 2000). Water samples were not preserved properly and not analyzed for concentration and stable carbon-isotopes of DIC. Previously however, the water had been repeatedly collected and analyzed, and the results gave relatively consistent DIC values for each facies (Fouke et al., 2000). We use these values as well as the reported $\delta^{13}\text{C}$ of DIC as references for this study (Table 2). The isotopic composition of dissolved CO_2 is estimated to be in a narrow range of -3.9‰ to -5.9‰ across these facies (Table 2). The $\delta^{13}\text{C}$ values of total biomass, however, show a much wider but

systematic variation along the temperature gradient, decreasing from -16.1‰ in the vent to -28.0‰ in the proximal slope followed by a slight increase to -23.5‰ in the distal slope (Table 2).

4.2. Lipid Profiles in Different Facies

4.2.1. Phospholipid fatty acids (PLFA)

PLFA in the vent facies are dominated by 20:1 ω 9c and 20:1 ω 9t, which collectively make up about one third (32.1%, average of samples A and B; same reporting below) of the total fatty acids (Table 3). The next most abundant fatty acid is 18:0 (22.2%) followed by 18:1 with three isomers, 18:1 ω 9c (8.2%), 18:1 ω 7c (7.6%), and 18:1 ω 7t (1.7%). Collectively, these isomers make up $\sim 17.4\%$ of total PLFA (Table 3). The universal fatty acid, 16:0, accounts for $\sim 10.0\%$ of total PLFA. Less abundant fatty acids include 20:0, cy19:0, and *iso*- or *anteiso*-15, -16 and -17 carbon chains, which are $<5\%$ each (Table 3).

The pond-2 and the proximal slope-2 (Fig. 1) have similar lipid profiles with 16:0 accounting for about half of total PLFA in both facies (Table 3). This is followed by 18:1 ω 9c and 18:1 ω 7c, which collectively account for $\sim 15.5\%$ of total PLFA in both facies. The most abundant fatty acid (20:1 ω 9) in the vent nearly disappears in these facies and a15:0, i17:0, a17:0, and cy19:0 also decrease to below 1.0% (Table 3). On the other hand, 16:1 ω 7c and 18:2 ω 6 appear in relatively high abundance (5.0–6.0%) with cy17:0 being a minor compound (1.5%) in the pond-2. In addition, the proximal slope-2 shows a sharp increase in 14:0, which accounts for $\sim 2.7\%$ of total PLFA (Table 3).

In the distal slope, 16:0 decreases to 30.0% of total PLFA but still remains the most abundant. The compound 18:0 continues to decrease to a value of 4.4%. On the other hand, 18:1 ω 9c remains relatively unchanged ($\sim 14.0\%$) compared to its abundances in the pond-2 or the proximal slope-2, but 18:1 ω 7c increases to a greater proportion of $\sim 11.0\%$. Collectively, 18:1 ω 9c and 18:1 ω 7c account for $\sim 25.0\%$ of total PLFA in the distal slope and are the most abundant in all four facies. Also increased in the distal slope are i15:0 ($\sim 7.0\%$), a15:0 ($\sim 1.3\%$), 16:1 ω 7c ($\sim 6.8\%$), i17:0 ($\sim 1.3\%$), 18:3 ω 3 ($\sim 5.4\%$), and 20:0–23:0 (1.0–2.2%) (Table 3). These changes may reflect shifts in microbial community structure from facies to facies.

4.2.2. Glycolipid fatty acids (GLFA)

In all facies GLFA have major fatty acids that are also found in PLFA (Table 4). However, the relative proportions of major fatty acids in glycolipids may be different from those in the phospholipids. For example, GLFA in the vent have a significantly greater proportion of 18:1 ω 9c (44.2%, sample B) and lower proportions of 20:1 ω 9c (4.8%, sample B) and 20:1 ω 9t (3.1%, sample B) than PLFA in the same facies (Table 4). In the pond-2 and the proximal slope-2, major GLFA such as 16:1 ω 7c, 16:0, 18:2 ω 6, 18:1 ω 9c and 18:0 have similar proportions as the PLFA (Table 4). In the distal slope, GLFA have greater proportions of 16:1 ω 7c (average 9.5% between samples A and B, same below) and 16:0 ($\sim 60.0\%$) but lower proportions of i15:0 (1.2%), 18:2 ω 6 ($\sim 3.4\%$), 18:1 ω 9c (7.1%), and 18:1 ω 7c (5.6%) than the PLFA (Table 4).

Table 3. Distribution of phospholipid fatty acids in microbial mats from different facies of Angel Terrace Spring AT-1.

Fatty acids (mol%)	Facies							
	Vent		Pond-2		Proximal slope-2		Distal slope	
	A	B	A	B	A	B	A	B
14:0	0.0	0.0	0.3	0.2	3.0	2.4	0.6	0.9
i15:0	2.3	1.5	1.8	1.4	1.8	2.0	6.5	7.5
a15:0	1.6	1.3	0.0	0.0	0.0	0.0	1.2	1.3
15:0	0.0	0.0	0.3	0.2	0.3	0.3	0.3	0.4
i16:0	1.5	1.2	2.8	2.3	2.2	2.8	0.6	0.6
16:1 ω 7c	0.0	0.0	6.2	5.2	3.7	4.6	6.8	6.7
16:1 ω 7t	0.0	0.0	0.7	0.6	0.0	0.0	0.3	0.4
16:1 ω 5c	0.0	0.0	0.1	0.0	0.2	0.4	1.0	1.1
cy16:0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0
16:0	9.4	10.0	54.7	58.7	58.2	50.7	29.6	30.3
i17:1 ω 7c	0.0	0.0	0.1	0.0	0.3	0.5	0.7	0.8
br17:1	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0
i17:0	3.9	4.6	0.5	0.4	0.3	0.4	1.4	1.2
a17:0	3.5	4.5	0.5	0.4	0.3	0.4	0.5	0.7
17:1 ω 6c	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.0
cy17:0	0.0	0.0	1.7	1.3	0.0	0.1	0.0	0.0
17:0	1.2	1.2	0.6	0.5	0.5	0.5	0.6	0.5
18:3 ω 6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3
i18:0	0.6	0.6	0.0	0.0	0.0	0.0	0.0	0.0
18:2 ω 6	0.0	0.0	5.5	4.3	5.8	7.9	6.1	6.8
18:3 ω 3	0.0	0.0	0.0	0.3	0.9	1.3	4.8	5.9
18:1 ω 9c	9.3	7.0	13.7	13.7	12.9	15.6	15.7	12.3
18:1 ω 7c	7.2	8.0	1.6	1.5	1.2	1.4	11.0	10.5
18:1 ω 7t	1.3	2.0	0.4	0.0	0.0	0.0	0.0	0.0
18:1 ω 5c	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2
18:0	19.6	24.7	7.8	8.4	7.9	7.5	4.2	4.4
10Me18:0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0
i19:0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
cy19:0	2.5	1.4	0.0	0.5	0.0	0.0	0.0	0.0
19:0	0.0	0.0	0.7	0.0	0.1	0.1	0.7	0.2
20:2 ω 6	0.0	0.0	0.0	0.0	0.2	0.5	0.8	1.4
20:1 ω 9c	25.5	18.6	0.0	0.0	0.1	0.2	0.0	0.5
20:1 ω 9t	8.6	11.5	0.0	0.0	0.0	0.0	0.3	0.4
20:0	1.9	2.0	0.0	0.1	0.1	0.2	1.1	1.0
22:0	0.0	0.0	0.0	0.0	0.0	0.0	2.3	1.9
23:0	0.0	0.0	0.0	0.0	0.0	0.0	2.7	1.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

4.3. Stable Isotope Compositions of PLFA and GLFA

Isotopic compositions of individual PLFA and GLFA were obtained only for those having concentrations high enough to be detected on the isotope-ratio-mass-spectrometer via the GC (Table 5, Table 6). Each sample was analyzed at least twice for the isotopic compositions of individual fatty acids. The average of the replicate runs is reported, which has a standard deviation less than 1.0‰ in most samples (Table 5, Table 6). Large standard deviations are associated with PLFA 18:0 (± 2.2 –4.6‰) in the distal slope (Table 5).

4.3.1. Isotopic compositions of PLFA

PLFA in the vent have $\delta^{13}\text{C}$ values ranging from $-18.7 \pm 5.6\%$ (average of samples A and B, $n = 2$; same reporting below unless otherwise noted) for 16:0 to $-6.3 \pm 1.6\%$ ($n = 2$) for $\Sigma 18:1\omega 9c/7c$. Collectively, the PLFA in the vent have an average value of $-12.4 \pm 5.2\%$ ($n = 10$, Fig. 3).

PLFA in the pond-2 and in the proximal slope-2 have $\delta^{13}\text{C}$

values with similar variations. For example, i15:0 is most enriched in ^{13}C and has $\delta^{13}\text{C}$ values of -26.7% in the pond-2 and $-26.8 \pm 0.3\%$ ($n = 2$) in the proximal slope-2 (Table 5). The second most ^{13}C -enriched fatty acid is i16:0, which has $\delta^{13}\text{C}$ values of -27.1% and $-28.6 \pm 0.2\%$ ($n = 2$) in these two facies, respectively. The fatty acid 16:1 ω 7c is mostly depleted in ^{13}C and has $\delta^{13}\text{C}$ values of $-35.4 \pm 0.4\%$ ($n = 2$) in the pond-2 and $-37.8 \pm 0.3\%$ ($n = 2$) in the proximal slope-2 (Table 5). Collectively, PLFA have average values of $-33.0 \pm 3.1\%$ ($n = 11$) in the pond-2 and $-33.7 \pm 3.8\%$ ($n = 16$) in the proximal slope-2 (Fig. 3). These average values are $\sim 21\%$ more negative than that in the vent (Fig. 3), suggesting a dramatic change in the mechanisms of carbon-isotope fractionation in these facies.

In the distal slope, isotopic compositions of individual PLFA become enriched in ^{13}C but not to the level of enrichment observed in the vent. The $\delta^{13}\text{C}$ ranges from $-18.4 \pm 0.1\%$ ($n = 2$) for i15:0 to $-27.0 \pm 0.2\%$ ($n = 2$) for 16:0. One exception is the isomers 18:1 ω 9c and 18:1 ω 7c, which have a

Table 4. Distribution of glycol-lipid fatty acids in microbial mats from different facies of Angel Terrace Spring AT-1.

Fatty acids (mol%)	Facies							
	Vent		Pond-2		Proximal slope-2		Distal slope	
	A ^a	B	A	B	A	B	A	B
14:0	0.0	0.0	0.8	0.0	7.1	6.7	1.3	1.6
i15:0	0.0	0.0	0.4	0.0	0.3	0.3	1.0	1.3
a15:0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
15:0	0.0	0.0	0.2	0.0	0.1	0.1	0.0	0.2
i16:0	0.0	0.0	0.5	0.0	0.4	0.4	0.0	0.2
16:1	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.5
16:1 ω 7c	0.0	0.0	8.9	5.9	6.4	6.0	10.2	8.8
16:1 ω 7t	0.0	0.0	0.3	0.0	0.1	0.1	0.0	0.4
16:0	0.0	11.9	62.0	72.1	52.4	49.7	61.3	58.6
i17:0	0.0	0.7	0.2	0.0	0.1	0.1	0.5	0.6
a17:0	0.0	1.3	0.0	0.0	0.0	0.1	0.0	0.1
Unknown	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2
17:0	0.0	1.1	0.3	0.0	0.2	0.2	0.4	0.5
18:2 ω 6	0.0	1.4	6.8	5.1	13.6	14.2	3.0	3.8
18:3 ω 3	0.0	0.0	0.4	0.0	1.4	1.8	3.5	4.0
18:1 ω 9c	0.0	44.2	12.4	12.3	12.7	12.5	9.3	4.9
18:1 ω 7c	0.0	9.7	1.0	0.0	0.9	1.1	5.8	5.4
18:1 ω 7t	0.0	1.7	0.0	0.0	0.1	0.0	0.0	0.1
18:1 ω 5c	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
18:0	0.0	18.5	5.8	4.6	2.7	3.1	3.6	3.5
10Me18:0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
19:2	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.2
br19:0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
br19:1	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
br19:0	0.0	0.0	0.0	0.0	0.0	2.3	0.0	0.1
20:2 ω 6	0.0	0.0	0.0	0.0	0.2	0.2	0.0	1.5
20:3 ω 3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
20:1 ω 9c	0.0	4.8	0.0	0.0	0.1	0.0	0.0	0.0
20:1 ω 9t	0.0	3.1	0.0	0.0	0.1	0.1	0.0	0.3
20:0	0.0	1.5	0.0	0.0	0.1	0.1	0.0	0.4
br20:0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.2
br21:0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0
22:2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
Unknown	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
22:1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
22:0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6
Unknown	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
23:0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
24:0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4
Total	0.0	100.0	100.0	100.0	99.6	100.0	100.0	99.4

^a Sample was lost.

composite $\delta^{13}\text{C}$ of -40.0% (Table 5). This value is suspiciously low compared with $\delta^{13}\text{C}$ values of other PLFA in the same facies. Because we didn't have a replicate measurement in sample A to confirm this value, it is excluded for further calculation. As a result, the overall PLFA have an average value of $-22.4 \pm 3.4\%$ ($n = 10$) in the distal slope (Fig. 3). This is approximately 11‰ enriched in ^{13}C relative to PLFA in the pond-2 and the proximal slope-2 and $\sim 10\%$ depleted in ^{13}C relative to PLFA in the vent (Fig. 3).

4.3.2. Isotopic compositions of GLFA

Fewer fatty acids in the glycolipids were available for isotopic analyses (Table 6). Nevertheless, the data show that most of the GLFA are depleted in ^{13}C relative to the same fatty acids in PLFA. In the vent, for example, the $\delta^{13}\text{C}$ values of individual GLFA are depleted by 7.6‰ (18:0) to 11.5‰ ($\Sigma 18:1\omega 9/7c$)

relative to the same fatty acids in PLFA (Table 5). Most of the GLFA in the proximal slope and the distal slope are also depleted in ^{13}C (0.1–8.2‰) relative to the PLFA. Exceptions are 18:2 ω 6 and 18:0 in sample B of the proximal slope-2, for which the GLFA is ~ 1 –3.3‰ enriched in ^{13}C relative to PLFA (Table 5, Table 6). In the pond-2, however, most of the GLFA are enriched in ^{13}C (2.8–6.0‰) relative to PLFA with the exception of 16:0 (Table 5, Table 6).

The average $\delta^{13}\text{C}$ values of GLFA vary from facies to facies in a manner similar to that of PLFA. For example, the average $\delta^{13}\text{C}$ of GLFA is $-19.6 \pm 3.4\%$ ($n = 3$) in the vent, $-30.4 \pm 4.7\%$ ($n = 8$) in the pond-2, $-36.9 \pm 2.8\%$ ($n = 12$) in the proximal slope-2, and $-27.9 \pm 3.1\%$ ($n = 13$) in the distal slope (Fig. 3). The isotopic variations of PLFA and GLFA are also consistent with that of total biomass among these facies (Fig. 3).

Table 5. Isotopic compositions of phospholipid fatty acids of microbial mats from different facies of Angel Terrace Spring AT-1.

Fatty acid	$\delta^{13}\text{C}$ (‰) ^a							
	Vent		Pond-2		Proximal slope-2		Distal slope	
	A	B	A	B	A	B	A	B
14:0					-35.2 (0.4)	-35.4 (0.1)		
i15:0				-26.7 (0.2)	-27.0 (0.2)	-26.6 (0.1)	-18.5 (0.2)	-18.3 (0.1)
i16:0				-27.1 (0.4)	-28.8 (0.2)	-28.5 (0.8)		
16:1 ω 7c			-35.1 (1.0)	-35.7 (0.1)	-37.7 (0.2)	-38.0 (0.1)	-19.6 (0.0)	-19.6 (0.5)
16:0	-22.7 (0.4)	-14.8 (0.3)	-34.7 (0.1)	-35.1 (0.0)	-36.3 (0.1)	-36.0 (0.1)	-26.8 (0.0)	-27.1 (0.1)
Σ i17:0,a17:0 ^b	-16.7	-14.8						
18:2 ω 6				-34.3 (0.3)	-34.6 (0.3)	-34.9	-22.8 (0.2)	-23.2
Σ 18:1 ω 9/7c ^b	-7.4	-5.2	-33.2	-33.8	-36.7	-35.0		-40.0
18:0	-13.8 (0.5)	-11.9 (1.1)	-33.1 (0.5)	-34.0 (0.1)	-34.7 (0.4)	-33.7	-26.0 (4.6)	-21.8 (2.2)
Σ 20:1 ω 9t/c ^b	-8.2	-8.7						

^a Average \pm 1 standard deviation (in parentheses) of two or three replicate runs. The $\delta^{13}\text{C}$ values of individual fatty acids were corrected for methylation (see text).

^b Baseline resolution was not possible; a composite $\delta^{13}\text{C}$ value was obtained for coeluting fatty acids by integrating individual $\delta^{13}\text{C}$ values using weight percentage of mass 44 [V] of each peak.

DISCUSSION

5.1. Lipid Profiles of Different Facies

Fouke et al. (2003) have provided a phylogenetic survey of bacterial species in each facies at Angel Terrace Spring AT-1 based on 16S rRNA gene sequences. The results show strong partitioning between dominant microbial species and types of depositional facies (Table 1). In particular, the vent is dominated by *Aquificales* (91% of total sequences). Gene sequences of the green sulfur bacteria (i.e., *Chlorobium*) are most abundant in the proximal slope (9%) and less than 1% in the vent; those of the green non-sulfur bacteria (i.e., *Chloroflexus*) range from 4% in the apron and channel to below detection in the distal slope (Table 1). The cyanobacteria have consistently higher proportions in the relatively low-temperature environments (the pond, the proximal slope and the distal slope) than in the high-temperature environments (the vent and the apron and channel) (Table 1). Except in the vent, the majority of gene sequences belong to other bacteria (Table 1), whose roles in these environments are not well defined. It must be understood,

however, that the relative proportion of sequences (clones) representing a given species or division of bacteria is not directly correlated to the actual size of the cellular population of that species or division.

Lipid biomarkers can provide valuable information about microbial structures in geological settings. Pioneering studies of microbial lipids in Yellowstone hot springs were led by Ward and colleagues, who performed extensive analyses of lipid biomarkers of cyanobacterial mats (Ward et al., 1985; Dobson et al., 1988; Ward et al., 1989; Zeng et al., 1992a,b). Zeng et al. (1992a) observe that the distribution of most of the major mat lipids in the Yellowstone hot springs is consistent with lipid profiles of microorganisms isolated from the mats. Jahnke et al. (2001) focus their study on the lipid biomarkers in the Octopus Spring hyperthermophilic community in Yellowstone National Park. Lipid extracts of the natural samples show fatty acid biomarkers dominated by cy21:0, 20:1 ω 9, i17:0, i19:0, and 18:0. The abundance of 20:1 ω 9, cy21:0, and 18:0 are characteristic of certain known *Aquificales* species (Jahnke et al., 2001). Other culture studies identify 18:1 ω 9/7 as additional

Table 6. Isotopic compositions of glycolipid fatty acids of microbial mats from different facies of Angel Terrace Spring AT-1.

Fatty acid	$\delta^{13}\text{C}$ (‰) ^a							
	Vent		Pond-2		Proximal slope-2		Distal slope	
	A	B	A	B	A	B	A	B
14:0			-20.4		-37.6 (0.3)	-36.7 (0.2)	-32.9	-29.6
i15:0							-21.2 (0.1)	-23.8 (0.5)
16:1 ω 7c			-32.1	-29.7	-41.1 (0.1)	-39.8 (0.2)	-27.0 (1.1)	-26.2 (0.0)
16:0		-22.6	-35.5	-35.3	-37.9 (0.1)	-37.4 (0.2)	-29.9 (0.2)	-29.3 (0.4)
18:2 ω 6					-35.9 (0.3)	-33.9 (0.2)	-31.0	-26.8 (0.3)
Σ 18:1 ω 9/7c ^b		-16.7	-30.3	-31.0	-38.7 (0.0)	-38.1 (0.0)		-30.5
18:0		-19.5	-28.5		-34.8 (0.2)	-30.4 (0.2)	-26.7	-27.4 (0.5)

^a Average \pm 1 standard deviation (in parentheses) of two or three replicate runs. The $\delta^{13}\text{C}$ values of individual fatty acids were corrected for methylation (see text).

^b Baseline resolution was not possible; a composite $\delta^{13}\text{C}$ value was obtained for co-eluting fatty acids by integrating individual $\delta^{13}\text{C}$ values using weight percentage of mass 44 [V] of each peak.




Figure 3. Average isotopic compositions of phospholipids fatty acids (PLFA), glycolipid fatty acids (GLFA), and total biomass in different facies at Angel Terrace Spring AT-1. The isotopic compositions of dissolved CO₂ calculated from total dissolved inorganic carbon (Table 2) are also shown.

major fatty acids instead of cy21:0 (Kawasumi et al., 1984; Nishihara et al., 1990; Stohr et al., 2001a; Zhang et al., 2002).

The above discussion provides a basis for understanding the biogeochemical dynamics of lipids in different facies at Spring AT-1. For example, the dominance of 20:1 ω 9, 18:0, and 18:1 ω 9/7 in the vent (Table 3, Table 4) is consistent with the distribution of the 16S rRNA gene sequences of *Aquificales* in the vent (Fouke et al., 2003) and consistent with lipid profiles of known *Aquificales* species (Kawasumi et al., 1984; Huber et al., 1998; Jahnke et al., 2001; Stohr et al., 2001a; Gotz et al., 2002; Zhang et al., 2002). However, *Aquificales* in pure cultures do not produce *iso*- or *anteiso*- fatty acids (Jahnke et al., 2001; Zhang et al., 2002). Thus the presence of *iso*- or *anteiso*-15:0 to -17:0 in the vent are likely derived from other bacterial species. These may include *Thermus aquaticus*, *Bacillus stearothermophilus* and other mat-inhabiting heterotrophs, which are known to produce *iso*- and *anteiso*-15:0 to 17:0 compounds as major fatty acids (Yao et al., 1970; Ray et al., 1971; Oshima and Miyagawa, 1974).

The similar lipid profiles between the pond-2 and the proximal slope-2 suggest that the relative proportions of extant microbial populations in these two facies may be similar. The dramatic decrease in 20:1 ω 9c/t in these facies suggests that the relative abundance of *Aquificales* is much lower than in the vent (Table 3, Table 4). On the other hand, 16:1 ω 7c and 18:2 ω 6 in both PLFA and GLFA begin to appear in significant abundance accompanied by increasing 16:0 (Table 3, Table 4), which are consistent with their presence in cyanobacteria or green non-sulfur bacteria (Fork et al., 1979; Ward et al., 1989; Zheng et al., 1992a; Summons et al., 1996).

In the distal slope, the microbial community is also likely dominated by cyanobacteria and green non-sulfur bacteria as demonstrated by high abundances of 16:1 ω 7c and 18:2 ω 6, and 18:1 ω 9c that are similar to those in the pond-2 and the proximal slope-2. The increase in 18:3 ω 3 in the distal slope may further

indicate the presence of cyanobacteria because 18:3 biomarkers have been reported as major fatty acids in cyanobacterium *Phormidium luridum* grown at temperatures similar to the distal slope environment (Summons et al., 1996). In addition, the increase in i15:0 may indicate an increasing population of heterotrophic bacteria growing at low temperatures.

Overall, the lipid profiles clearly reflect the changing microbial communities from the vent to the pond and the proximal slope and to the distal slope. Temperature appears to be the major controlling factor for the dominant microbial species in these environments. For example, most *Aquificales* can grow at temperatures above 70°C (Kawasumi et al., 1984; Huber et al., 1998, 2002; Reysenbach et al., 2000; Stohr et al., 2001a; Gotz et al., 2002); whereas, cyanobacteria, green sulfur bacteria, green non-sulfur bacteria, and *Hydrogenophilus hirschii* Yel5a grow at temperatures below 60–70°C (Meeks and Castenholz, 1971; Bauld and Brock, 1973; Ward et al., 1998; Miller and Castenholz, 2000; Stohr et al., 2001b). Algae and other eukaryotes are normally found at temperatures below 40–55°C (Castenholz, 1984). On the other hand, fatty acid compositions of individual cells can also vary under different growth temperatures (Ray et al., 1971; Fork et al., 1979; Jahnke, 1992; Könneke and Widdel, 2003). In particular, variations in temperature can induce changes in unsaturation of fatty acids necessary for proper function of cell membranes (Murata et al., 1992; Murata and Wada, 1995; Könneke and Widdel, 2003). Overall, results of this study support similar patterns of temperature distribution of microbial species in hot springs that were determined using molecular DNA techniques (Ward et al., 1998; Miller and Castenholz, 2000; Fouke et al., 2003).

5.2. Carbon Isotopic Compositions of Fatty Acids

The estimated isotopic compositions of dissolved CO₂ (Table 2) show small variation ($\leq 2.0\text{‰}$) with decreasing DIC from the vent (192 ppm) to the distal slope (75 ppm); whereas, the isotopic compositions of total biomass show a large variation decreasing from -16.1‰ in the vent to -23.5‰ in the distal slope (Table 2). These results suggest that the abundance and isotopic compositions of CO₂ or DIC do not seem to have major influence on the changing isotopic compositions of organic matter in these environments. One exception may be the distal slope, where degassing results in 3.0‰ enrichment in ¹³C_{DIC} relative to DIC in the upper proximal slope. This might have contributed to the 4.5‰ increase in ¹³C_{Biomass} in the distal slope relative to biomass in the proximal slope (Table 2).

Isotopic compositions of PLFA and GLFA are consistent with those of total biomass between different facies (Fig. 3). This suggests that in each facies fatty acids and biomass may reflect the same mechanism of isotope fractionation. Different facies, however, appear to be associated with different mechanisms of isotope fractionation as demonstrated by the plot of $\epsilon_{\text{FA-Bio}}$ vs. $\epsilon_{\text{Bio-CO}_2}$ (Fig. 4). In this plot, the trend of fractionations for different facies is parallel to that for pure cultures, among which the cyanobacteria represent fractionations for the Calvin cycle (lower end-member), the *Aquificales* represent fractionations for the rTCA cycle (upper end-member), and the green non-sulfur bacteria represent the 3-HP pathway between the two end-members (Fig. 4). Such a comparison indicates that CO₂ fixation in the vent may be dominated by rTCA; whereas,




Figure 4. Isotope fractionations between fatty acids (FA) and total biomass (Bio) vs. fractionations between biomass (Bio) and CO₂ for pure cultures (open circles) and for environmental samples from different facies at AT-1 (solid circles). For pure cultures, Calvin represents cyanobacteria (*Synechocystis* UTEX 2470 and *Phormidium luridum*), rTCA represents *Aquificales* (*Persephonella marina* and *Thermocrinis ruber*), and 3-HP represents green non-sulfur bacteria (e.g., *Chloroflexus aurantiacus*) (Table 7).

CO₂ fixation in the pond-2 and the proximal slope-2 may be dominated by the Calvin cycle. CO₂ fixation in the distal slope is consistent with the 3-HP pathway; however, the fractionation can also represent a mixture of the two end-members.

Among the proposed mechanisms of CO₂ fixation, the rTCA cycle can be unambiguously identified in the vent because of the dominance of 20:1 ω 9 and 18:1 ω 9/7 biomarkers that are specific for *Aquificales*. This, coincidentally, also corresponds to the large proportion of gene sequences of *Aquificales* in the vent (Fouke et al., 2003).

In the pond-2 and the proximal slope-2, it may be difficult to distinguish the relative contributions of cyanobacteria, green non-sulfur bacteria, and/or green sulfur bacteria to the biomass production because these organisms may closely interact with each other. For example, it has been observed that *Chloroflexus* spp. of the green non-sulfur bacteria, are closely associated with cyanobacteria and uses their organic byproducts for photoheterotrophic growth. When this happens, the mixture of these species shows the isotopic composition of the cyanobacteria (Estep, 1984). If that is the case in the pond-2 and the proximal slope-2, we would expect $\delta^{13}\text{C}$ of *Chloroflexus* to be similar to that of cyanobacteria, thus underestimating their contribution to the community biomass.

In the distal slope, the fractionation is consistent with the 3-HP pathway, which may be used by the green non-sulfur bacteria such as *Chloroflexus* (Table 7). However, gene sequencing shows a lack of green non-sulfur bacteria in the distal slope (Fouke et al., 2003). Thus it is not clear whether *Chloroflexus* contributes significantly to biomass production in the distal slope. On the other hand, the fractionations can be explained by a mixture of cyanobacteria with more negative fractionations and green sulfur bacteria (i.e., *C. limicola*) with more positive fractionations (Table 7). Both groups are present in the distal slope based on gene sequencing (Fouke et al., 2003).

Large differences in isotopic compositions also occur between individual fatty acids within each facies (Table 5, Table

Table 7. Carbon isotopic fractionations of microbial species associated with known biosynthetic pathways.

Microorganisms	$\delta^{13}\text{C}_{\text{CO}_2}$ (‰)	$\delta^{13}\text{C}_{\text{biomass}}$ (‰)	$\delta^{13}\text{C}_{\text{fatty acids}}$ (‰)	$\epsilon_{\text{FA-BIO}}^{\text{a}}$ (‰)	$\epsilon_{\text{BIO-CO}_2}^{\text{a}}$ (‰)	$\epsilon_{\text{FA-CO}_2}^{\text{a}}$ (‰)	Biosyn. pathway	Ref. ^b
Cyanobacteria								
<i>Synechocystis</i> UTEX 2470	-9.9	-30.7	-39.6	-9.2	-21.0	-30.0	Calvin	1
<i>Synechococcus</i> sp.	-7.8 to -16.9	-26.1 to -33.1			-17.0 \pm 1.2		Calvin	2
<i>Phormidium luridum</i>	-7.8 to -39.9	-18.4 to -60.1	-28.7 to -66.4	-9.1 \pm 1.5	-14.3 \pm 4.4	-23.3 \pm 2.9	Calvin	3
Algae								
<i>Tetraedron minimum</i>				-5.0			Calvin	4
<i>Porosira glacialis</i>	-28.3 to -37.9	-34.2 to -58.8			-14.1 \pm 5.2		Calvin	2
<i>Aquificales</i>								
<i>Persephonella marina</i>	-40.8	-45.6	-42.9	2.8	-5.0	-2.2	r-TCA	5
<i>Thermocrinis ruber</i>	-27.4	-30.7	-28.4	2.4	-3.4	-1.0	r-TCA	6
Green sulfur bacterium								
<i>Chlorobium limicola</i>		-26.3		12.9			r-TCA	7
<i>Chlorobium thiosulfatophilum</i>	-16.5 ^c	-28.7			-12.2		r-TCA	8
Green non-sulfur bacterium								
<i>Chloroflexus aurantiacus</i>	-41.7	-48.9	-49.6	-0.7	-7.6	-8.4	3-HP ^d	9

^a Fractionation between two substances. $\epsilon_{\text{A-B}} = ((\delta^{13}\text{C}_{\text{A}} + 1000)/(\delta^{13}\text{C}_{\text{B}} + 1000) - 1) \times 1000$. FA = total fatty acids; BIO = biomass or total organic carbon. In cases of average \pm 1 standard deviation; n = 5–12.

^b 1. Sakata et al. (1997); 2. Popp et al. (1998); 3. Summons et al. (1996); 4. Schouten et al. (1998); 5. Zhang et al. (2002); 6. Jahnke et al. (2001); 7. van der Meer et al. (1998); 8. Sirevåg et al. (1977); 9. van der Meer et al. (2001).

^c Calculated from measurement of $^{13}\text{C}_{\text{HCO}_3^-}$ using a fractionation factor of 7.3‰ between dissolved CO₂ and bicarbonate as used by Sirevåg et al. (1977).

^d Although the 3-HP pathway uses HCO₃⁻ rather than CO₂ as a substrate for cell growth, CO₂ is used for calculation in order to compare with other pathways using CO₂ for cell growth.

6). These variations may reflect: 1) different biosynthetic pathways of different microorganisms, 2) different carbon substrates used by the same microorganism, or 3) variation within the same microorganism. Factor one has been discussed above. Factors 2 and 3 cannot be determined using the existing data. However, one example is given by Jahnke et al. (2001) who show that isotopic compositions of lipids of *Thermocrinis ruber* (rTCA cycle) were $\sim 10\text{‰}$ lighter when grown heterotrophically on formate (-23.3‰) than grown autotrophically on CO_2 (-27.4‰), even though the isotopic compositions of CO_2 and formate are similar. Nevertheless, isotopic compositions of fatty acid biomarkers provide valuable information about changing microbial community structure and biosynthetic pathways from the vent to the pond and the proximal slope to the distal slope at Spring AT-1.

5.3. Geological and Evolutionary Implications

Previous studies of travertine deposits have focused on the inorganic chemistry of the spring water and carbonate mineralogy. While travertine crystal fabrics are strongly influenced by microbes, which may provide a substrate for crystal nucleation, the chemical byproducts of microbial respiration and photosynthesis are not well expressed in the travertine crystal geochemistry. This study integrates lipid biomarkers and stable isotopes in the context of microbial communities, water chemistry, and varying facies of carbonate precipitation, which provide substantial insights into the relationships between microbial population dynamics and the changing environment in travertine deposits.

Studies of hot springs in other locations around the world show lipid distributions that are consistent with our observations at Mammoth Hot Springs. For example, hot springs in Iceland and New Zealand have lipid biomarkers consistent with the dominance of cyanobacterial mats (Robinson and Eglinton, 1990; Shiea et al., 1990, 1991). A variety of hot springs and hydrothermal vents harbor *Aquificales*, which all have the characteristic fatty acids of 20:1 ω 9, 18:0, 18:1 ω 9/7, or cy21:0 (Kawasumi et al., 1984; Huber et al., 1998; Jahnke et al., 2001; Stohr et al., 2001a; Gotz et al., 2002; Zhang et al., 2002). The latter biomarkers are rare in other bacterial species and may serve as taxonomic biomarkers for *Aquificales* species (Kawasumi et al., 1984; Jahnke et al., 2001). This study has enriched our knowledge about lipid biomarker and isotopic signatures of major microbial populations in geological settings.

Aquificales, Cyanobacteria, green sulfur bacteria, and green non-sulfur bacteria have been frequently observed in hot springs in the Yellowstone National Park (Ward et al., 1984, 1997, 1998; Stahl et al., 1985; Reysenbach et al., 1994; Hugenholtz et al., 1998; Eder and Huber, 2002). These species have also been found in hot springs in other regions of the world (Cohen, 1984; Robinson and Eglinton, 1990; Eder and Huber, 2002; Nakagawa and Fukui, 2002). They are of particular interest for studies of microbial evolution. For example, the *Aquificales* represents one of the deepest lineages of the Bacteria in the phylogenetic tree (Reysenbach et al., 1994). Green sulfur and green non-sulfur bacteria are anaerobic phototrophs and might have evolved before the atmosphere was oxygenated (Oyaizu et al., 1987; Gupta et al., 1999). The cyanobacteria, on the other hand, may provide clues about the

early stages of arising oxygen levels on earth (Summons et al., 1996). Recent advances in molecular microbiology and compound-specific isotope signatures will shed new light and provide more precise information about biogeochemical processes mediated by these organisms in modern systems, which help better understand the evolution of life in ancient environments.

SUMMARY AND CONCLUSIONS

Phospholipid fatty acids (PLFA), glycolipid fatty acids (GLFA), and their carbon isotope compositions were determined for microbial mats and rock materials from four different facies that comprise the modern travertine deposits in the Spring AT-1 outflow drainage system of Mammoth Hot Springs in Yellowstone National Park. In the vent (71°C), PLFA and GLFA are dominated by 20:1 ω 9, 18:0, and 18:1 ω 9/7, which are consistent with biomarkers observed in *Aquificales*; isotope fractionations between fatty acids and biomass are consistent with rTCA cycle used by *Aquificales* for CO_2 fixation. In the pond (51°C) and the proximal slope (47°C), characteristic biomarkers include 16:1 ω 7c, 18:1 ω 9/7, and 18:2 ω 6, which may represent cyanobacteria, green non-sulfur bacteria, and/or green sulfur bacteria; isotope fractionations are consistent with the Calvin cycle being a potential mechanism for CO_2 fixation. Lipid biomarkers in the distal slope (24°C) also indicate the predominance of cyanobacteria, green non-sulfur bacteria and/or green sulfur bacteria; isotope fractionations, however, are consistent with the 3-HP pathway or a mixture of the Calvin cycle or rTCA cycle.

Results of this study show that lipid biomarkers and their isotopic compositions are only slightly affected by abundance and isotopic compositions of dissolved inorganic carbon in different facies. Instead, lipid biomarkers mostly reflect the changing microbial communities in different environments with temperature being the dominating factor; the isotopic compositions of biomass and fatty-acid biomarkers, on the other hand, provide a valuable insight into the mechanisms of CO_2 fixation by the extant microbial communities in the changing environments.

Stetter, 1994

Acknowledgments—Qi Ye and James Cantu provided technique support for lipid extraction. We thank Dr. Lesley Warren for including our paper in this special volume on Microbial Geochemistry. Comments from three anonymous reviewers and Editor Dr. Jan Amend significantly enhanced the quality of the manuscript. This research was partially supported by the Environmental Remediation Sciences Division of the Office of Biologic and Environmental Research, U.S. Department of Energy through the Financial Assistant Award no. DE-FC09-96SR18546 to the University of Georgia Research Foundation (CLZ). Additional partial supports were from the National Science Foundation Biocomplexity in the Environment Coupled Biogeochemical Cycles Program (BWF), the National Science Foundation Geosciences Postdoctoral Research Fellowship Program (GTB), the Petroleum Research Fund of the American Chemical Society Starter Grant Program (BWF), and the University of Illinois Urbana-Champaign Critical Research Initiative (BWF).

REFERENCES

- Abrajano T. A. J., Murphy D. E., Fang J., Comet P., and Brooks J. M. (1994) $^{13}\text{C}/^{12}\text{C}$ ratios in individual fatty acids of marine mytilids with and without bacterial symbionts. *Org. Geochem.* **21**, 611–617.

- Allen E. T. and Day A. L. (1935) Hot springs of the Yellowstone National Park pp. 525. Carnegie Institution of Washington.
- Bargar K. E. (1978) Geology and thermal history of Mammoth Hot Springs, Yellowstone National Park, Wyoming. *U. S. Geological Survey Bulletin* **1444**, 1–54.
- Barns S. M. and Nierzwick-Bauer S. (1997) Microbial diversity in modern subsurface, ocean, surface environments. In *Reviews in Mineralogy*, Vol. 35, pp. 35–79. Mineralogical Society of America.
- Bauld J. and Brock T. D. (1973) Ecological studies of *Chloroflexus*, a gliding photosynthetic bacterium. *Arch. Mikrobiol.* **92**, 267–284.
- Beh M., Strauss G., Huber R., Stetter K. O., and Fuchs G. (1993) Enzymes of the reductive citric acid cycle in the autotrophic eubacterium *Aquifex pyrophilus* and in the archaeobacterium *Thermoproteus neutrophilus*. *Arch. Mikrobiol.* **160**, 306–311.
- Brassell S. C. (1992) Biomarkers in recent and ancient sediments: the importance of the diagenetic continuum. In *Organic matter—productivity, accumulation, and preservation in recent and ancient sediments* (eds. J. K. Whelan and J. W. Farrington), pp. 339–367. Columbia University Press.
- Castenholz R. W. (1984) Composition of hot spring microbial mats: A summary. In *Microbial Mats: Stromatolites* (eds. Y. Cohen, R. W. Castenholz, and H. O. Halvorson), pp. 101–119. Alan R. Liss, Inc.
- Cohen Y. (1984) The Solar Lake cyanobacterial mats: strategies of photosynthetic life under sulfide. In *Microbial Mats: Stromatolites* (eds. Y. Cohen, R. W. Castenholz, and H. O. Halvorson), pp. 133–148. Alan R. Liss, Inc.
- De Rosa M., Gambacorta A., Huber R., Lanzotti V., Nicolaus B., Stetter K. O. and Trincone A. (1988) Lipid structures in *Thermotoga maritima*. *J. Chem. Soc. Chem. Commun.* 1300–1301.
- Dobson G., Ward D. M., Robinson N., and Eglinton G. (1988) Biogeochemistry of hot spring environments: extractable lipids of a cyanobacterial mat. *Chem. Geol.* **68**, 155–179.
- Eder W. and Huber R. (2002) New isolates and physiological properties of the *Aquificales* and description of *Thermocrinis albus* sp nov. *Extremophiles* **6**, 309–318.
- Estep M. L. (1984) Carbon and hydrogen isotopic compositions of algae and bacteria from hydrothermal environments, Yellowstone National Park. *Geochim. Cosmochim. Acta* **48**, 591–599.
- Farmer J. D. and Des Marais D. J. (1994) Biological versus inorganic processes in stromatolite morphogenesis: Observations from mineralizing sedimentary systems. In *Microbial Mats: Structure, Development, and Environmental Significance: NATO ASI Series in Ecological Sciences*, Vol. G35 (eds. L. J. Stal and P. Caumette), pp. 61–68. Springer-Verlag.
- Fork D. C., Murata N., and Sato N. (1979) Effect of growth temperature on the lipid and fatty acid composition and the dependence on temperature of light-induced redox reactions of cytochrome *f* and of light energy redistribution in the thermophilic blue-green alga *Synechococcus lividus*. *Plant Physiol.* **63**, 524–530.
- Fouke B. W., Farmer J. D., Des Marais D. J., Pratt L., Sturchio N. C., Burns P. C. and Discipulo M. K. (2000) Depositional facies and aqueous-solid geochemistry of travertine-depositing hot springs (Angel Terrace, Mammoth Hot Springs, Yellowstone National Park, U.S.A.). *J. Sed. Res.* **70**, 565–585.
- Fouke B. W., Bonheyo G. T., Sanzenbacher B., and Frias-Lopez J. (2003) Partitioning of bacterial communities between travertine depositional facies at Mammoth Hot Springs, Yellowstone National Park, USA. *Can. J. Earth. Sci.* **40**, 1531–1548.
- Freeman K. H. (2001) Isotopic biogeochemistry of marine organic carbon. In *Stable Isotope Geochemistry*, Vol. 43 (eds. J. W. Valley and D. R. Cole), pp. 579–605. The Mineralogical Society of America.
- Friedman I. (1970) Some investigations of the deposition of travertine from hot springs: I. The isotope chemistry of a travertine-depositing spring. *Geochim. Cosmochim. Acta* **34**, 1303–1315.
- Fuchs G. (1989) Alternative pathways of autotrophic CO₂ fixation. In *Autotrophic Bacteria* (eds. H. G. Schlegel and B. Bowien), pp. 365–382. Science Tech Publishers.
- Gotz D., Banta A., Beveridge T. J., Rushdi A. I., Simoneit B. R. T., and Reysenbach A. (2002) *Persephonella marina* gen. nov., sp nov and *Persephonella guaymasensis* sp nov., two novel, thermophilic, hydrogen-oxidizing microaerophiles from deep-sea hydrothermal vents. *Intl. J. Syst. Evol. Microbiol.* **52**, 1349–1359.
- Guckert J. B., Antworth C. B., Nichols P. D., and White D. C. (1985) Phospholipid ester-linked fatty acid profiles as reproducible assays for changes in prokaryotic community structure of estuarine sediments. *FEMS Microbiol. Ecol.* **31**, 147–158.
- Gupta R. S., Mukhtar T., and Singh B. (1999) Evolutionary relationships among photosynthetic prokaryotes (*Heliobacterium chlorum*, *Chloroflexus aurantiacus*, cyanobacteria, *Chlorobium tepidum* and proteobacteria): implications regarding the origin of photosynthesis. *Molecul. Microbiol.* **32**, 893–906.
- Hayes J. M. (2001) Fractionation of carbon and hydrogen isotopes in biosynthetic processes. In *Stable Isotope Geochemistry*, Vol. 43 (eds. J. W. Valley and D. R. Cole), pp. 225–277. The Mineralogical Society of America.
- Huber R., Wilharm T., Huber D., Trincone A., Burgraf S., König H., Rachel R., Rochinger I., Fricke H., and Stetter K. O. (1992) *Aquifex pyrophilus* gen. nov. sp. nov., represents a novel group of marine hyperthermophilic hydrogen-oxidizing bacteria. *System. Appl. Microbiol.* **15**, 340–351.
- Huber R., Eder W., Heldwein S., Wanner G., Huber H., Rachel R., and Stetter K. O. (1998) *Thermocrinis ruber* gen. nov., sp. nov., a pink-filament-forming hyperthermophilic bacterium isolated from Yellowstone National Park. *Appl. Environ. Microbiol.* **64**, 3576–3583.
- Huber H., Diller S., Horn C. and Rachel R. (2002) *Thermovibrio ruber* gen. nov., sp nov., an extremely thermophilic, chemolithoautotrophic, nitrate-reducing bacterium that forms a deep branch within the phylum *Aquificae*. *Intl. J. Syst. Evol. Microbiol.* **52**, 1859–1865.
- Hugenholtz P., Pitulle C., Hershberger K. L., and Pace N. R. (1998) Novel division level bacterial diversity in a Yellowstone hot spring. *J. Bacteriol.* **180**, 366–376.
- Jahnke L. L. (1992) The effects of growth temperature on the methyl sterol and phospholipid fatty-acid composition of *Methylococcus capsulatus* (Bath). *FEMS Microbiol. Lett.* **93**, 209–212.
- Jahnke L. L., Eder W., Huber R., Hope J. M., Hinrichs K. U., Hayes J. M., Des Marais D. J., Cady S. L. and Summons R. E. (2001) Signature lipids and stable carbon isotope analyses of octopus spring hyperthermophilic communities compared with those of *Aquificales* representatives. *Appl. Environ. Microbiol.* **67**, 5179–5189.
- Karl D. M. (1995) Ecology of free-living, hydrothermal vent microbial communities. In *The Microbiology of Deep-Sea Hydrothermal Vents* (ed. D. M. Karl), pp. 35–124. CRC Press.
- Kawasumi T., Igarashi Y., Kodama T., and Minoda Y. (1984) *Hydrogenobacter thermophilus* gen. nov., sp. nov., an extremely thermophilic, aerobic, hydrogen-oxidizing bacterium. *Int. J. System. Bacteriol.* **34**, 5–10.
- Könneke M. and Widdel F. (2003) Effect of growth temperature on cellular fatty acids in sulphate-reducing bacteria. *Environ. Microbiol.* **5**, 1064–1070.
- Langworthy T. A., Holzer G., Zeikus J. G., and Tornabene T. G. (1983) *Iso-* and *anteiso-*branched glycerol diethers of the thermophilic anaerobes *Thermodesulfobacterium commune*. *System. Appl. Microbiol.* **4**, 1–17.
- Londry K. L., Jahnke L. L. and Des Marais D. J. (2004) Stable carbon isotope ratios of lipid biomarkers of sulfate-reducing bacteria. *Appl. Environ. Microbiol.* **70**, 745–751.
- Machenzie A. S., Brassell S. C., Eglinton G., and Maxwell J. R. (1982) Chemical fossils—the geological fate of steroids. *Science* **217**, 491–504.
- Madigan M. T., Takigiku R., Lee R. G., Gest H., and Hayes J. M. (1989) Carbon isotope fractionation by thermophilic phototrophic sulfur bacteria: Evidence for autotrophic growth in natural populations. *Appl. Environ. Microbiol.* **55**, 639–644.
- Meeks J. C. and Castenholz R. W. (1971) Growth and photosynthesis in an extreme thermophile *Synechococcus lividus* (Cyanophyta). *Arch. Mikrobiol.* **78**, 25–41.
- Menendez C., Bauer Z., Huber H., Gadon N., Stetter K. O., and Fuchs G. (1999) Presence of acetyl-coenzyme A (CoA) carboxylase and propionyl-CoA carboxylase in autotrophic Crenarchaeota and indication for operation of a 3-hydroxypropionate cycle in autotrophic carbon fixation. *J. Bacteriol.* **181**, 1088–1098.
- Miller S. R. and Castenholz R. W. (2000) Evolution of thermotolerance in hot spring cyanobacteria of the genus *Synechococcus*. *Appl. Environ. Microbiol.* **66**, 4222–4229.

- Mook W. G., Brommerson J. C., and Staverman W. H. (1974) Carbon isotope fractionation between dissolved bicarbonate and gaseous carbon dioxide. *Earth Planet. Sci. Lett.* **22**, 169–176.
- Murata N. and Wada H. (1995) Acyl-lipid desaturases and their importance in the tolerance and acclimatization to cold of cyanobacteria. *Biochem. J.* **308**, 1–8.
- Murata N., Wada H., and Gombos Z. (1992) Modes of fatty-acid desaturation in cyanobacteria. *Plant Cell Physiol.* **33**, 933–941.
- Nakagawa T. and Hukui M. (2002) Phylogenetic characterization of microbial mats and streamers from a Japanese alkaline hot spring with a thermal gradient. *J. Gen. Appl. Microbiol.* **48**, 211–222.
- Nichols P. D., Guckert J. B., and White D. C. (1986) Determination of monounsaturated fatty acid double-bond position and geometry for microbial monocultures and complex microbial consortia by capillary GC/MS of their dimethyl disulfide adducts. *J. Microbiol. Meth.* **5**, 49–55.
- Nishihara H., Igarashi Y., and Kodama T. (1990) A new isolate of *Hydrogenobacter*, an obligately chemolithoautotrophic, thermophilic, halophilic and aerobic hydrogen-oxidizing bacterium from seaside saline hot-spring. *Arch. Microbiol.* **153**, 294–298.
- O'Leary M. H. (1988) Carbon isotopes in photosynthesis. *BioScience* **38**, 328–329.
- Oshima M. and Miyagawa A. (1974) Comparative studies on the fatty acid composition of moderately and extremely thermophilic bacteria. *Lipids* **9**, 476–480.
- Oyaizu H., Debrunner-Vossbrinck B., Mandelco L., Studier J. A., and Woese C. R. (1987) The green-sulfur bacteria: a deep branching in the eubacterial line of decent. *Syst. Appl. Microbiol.* **9**, 47–53.
- Popp B. N., Dore J. E., Hanson K. L., Wakeham S. G., Laws E. A., and Bidigare R. R. (1998) Effect of phytoplankton cell geometry on carbon isotopic fractionation. *Geochim. Cosmochim. Acta* **62**, 69–77.
- Preuß A., Schauder R., and Fuchs G. (1989) Carbon isotope fractionation by autotrophic bacteria with three different CO₂ fixation pathways. *J. Biosci.* **44C**, 397–402.
- Quandt L., Gottschalk G., Ziegler H., and Stichler W. (1977) Isotopic discrimination by photosynthetic bacteria. *FEMS Microbiol. Letts.* **1**, 125–128.
- Ratledge C. and Wilkinson S. G. (1988) *Microbial Lipids*, Vol. 1. Academic Press.
- Ray P. H., White D. C., and Brock T. D. (1971) Effect of temperature on the fatty acid composition of *Thermus aquaticus*. *J. Bacteriol.* **106**, 25–30.
- Reysenbach A. L., Wickham G. S., and Pace N. R. (1994) Phylogenetic analysis of the hyperthermophilic pink filament community in Octopus Spring, Yellowstone National Park. *Appl. Environ. Microbiol.* **60**, 2113–2119.
- Reysenbach A. L., Ehringer M., and Hershberger K. (2000) Microbial diversity at 83°C in Calcite Springs, Yellowstone National Park: Another environment where the *Aquificales* and “Korarchaeota” co-exist. *Extremophiles* **4**, 61–67.
- Ringelberg D. B., Davis J. D., Smith G. A., Pfiffner S. M., Nichols P. D., Nickels J. S., Henson J. M., Wilson J. T., Yates M., Campbell D. H., Read H. W., Stocksdale T. T., and White D. C. (1989) Validation of signature polarlipid fatty acid biomarkers for alkane-utilizing bacteria in soils and subsurface aquifer materials. *FEMS Microbiol. Ecol.* **62**, 39–50.
- Robinson N. and Eglinton G. (1990) Lipid chemistry of Icelandic hot spring microbial mats. *Org. Geochem.* **15**, 291–298.
- Sakata S., Hayes J. M., R. M. A., Evans R. A., Leckrone K. J. and Togasaki R. K. (1997) Carbon isotopic fractionation associated with lipid biosynthesis by a cyanobacterium: relevance for interpretation of biomarker records. *Geochim. Cosmochim. Acta* **61**, 5379–5389.
- Schouten S., Breteler W. C. M. K., Blokker P., Schogt N., Rijpstra W. I. C., Grice K., Bass M. and Sinninghe Damsté J. S. (1998) Biosynthetic effects on the stable carbon isotopic compositions of algal lipids: Implications for deciphering the carbon isotopic biomarker record. *Geochim. Cosmochim. Acta* **62**, 1397–1406.
- Shiea J., Brassell S. C., and Ward D. M. (1990) Mid-chain branched mono- and dimethyl alkanes in hot spring cyanobacterial mats: a direct biogenic source for branched alkanes in ancient sediments? *Org. Geochem.* **15**, 223–231.
- Shiea J., Brassell S. C., and Ward D. M. (1991) Comparative analysis of extractable lipids in hot spring microbial mats and their component photosynthetic bacteria. *Org. Geochem.* **17**, 309–319.
- Simoneit B. R. T. (1986) Cyclic terpenoids of the geosphere. In *Biological Markers in the Sedimentary Record* (ed. R. B. Johns), pp. 43–99. Elsevier Science Publishers.
- Simoneit B. R. T. (2002) Molecular indicators (biomarkers) of past life. *Anatomical Rec.* **268**, 186–195.
- Sirevåg R., Buchanan B. B., Berry J. A., and Troughton J. H. (1977) Mechanism of CO₂ fixation in bacterial photosynthesis studied by the carbon isotope fractionation technique. *Arch. Microbiol.* **112**, 35–38.
- Sorey M. L. and Colvard E. M. (1997) Hydrologic investigations in the Mammoth Corridor, Yellowstone National Park and vicinity, USA. *Geothermics* **26**, 221–249.
- Stahl D. A., Lane D. J., Olsen G. J., and Pace N. R. (1985) Characterization of a Yellowstone hot spring microbial community by 5S rRNA sequences. *Appl. Environ. Microbiol.* **49**, 1379–1384.
- Stohr R., Waberski A., Liesack W., Volker H., Wehmeyer U., and Thomm M. (2001a) *Hydrogenophilus hirschii* sp. nov., a novel thermophilic hydrogen-oxidizing beta-proteobacterium isolated from Yellowstone National Park. *Intl. J. Syst. Evol. Microbiol.* **51**, 481–488.
- Stohr R., Waberski A., Volker H., Tindall B. J. and Thomm M. (2001b) *Hydrothermus marinus* gen. nov., sp. nov., a novel thermophilic hydrogen-oxidizing bacterium, recognition of *Calderobacterium hydrogenophilum* as a member of the genus *Hydrogenobacter* and proposal of the reclassification of *Hydrogenobacter acidophilus* as *Hydrogenobaculum acidophilum* gen. nov., comb. nov., in the phylum ‘*Hydrogenobacter/Aquifex*’. *Intl. J. Syst. Evol. Microbiol.* **51**, 1853–1862.
- Strauss G. and Fuchs G. (1993) Enzymes of a novel autotrophic CO₂ fixation pathway in the phototrophic bacterium *Chloroflexus aurantiacus*, the 3-hydroxypropionate cycle. *Eur. J. Biochem.* **215**, 633–643.
- Summons R. E., Jahnke L. L., and Simoneit R. T. (1996) Lipid biomarkers for bacterial ecosystems: Studies of cultured organisms, hydrothermal environments and ancient sediments. In *Evolution of Hydrothermal Ecosystems on Earth (and Mars?)* (eds. G. R. Bock and J. A. Goode), pp. 174–194. Wiley.
- van der Meer M. T. J., Schouten S., and Sinninghe Damsté J. S. (1998) The effect of the reversed tricarboxylic-acid cycle on the ¹³C contents of bacterial lipids. *Org. Geochem.* **28**, 527–533.
- van der Meer M. T. J., Schouten S., van Dongen B. E., Rijpstra W. I. C., Fuchs G., Sinninghe Damsté J. S., de Leeuw J. W., and Ward D. M. (2001) Biosynthetic controls on the ¹³C contents of organic components in the phototrophic bacterium *Chloroflexus aurantiacus*. *J. Biol. Chem.* **276**, 10971–10976.
- Vorholt J., Kunow J., Stetter K. O., and Thauer R. K. (1995) Enzymes and coenzymes of the carbon monoxide dehydrogenase pathway for autotrophic CO₂ fixation in *Archaeoglobus lithotrophicus* and the lack of carbon monoxide dehydrogenase in the heterotrophic *A. profundus*. *Arch. Microbiol.* **163**, 112–118.
- Ward D. M., Beck E., Revsbech N. P., Sandbeck K. A., and Winfrey M. R. (1984) Decomposition of hot spring microbial mats. In *Microbial Mats: Stromatolites* (eds. Y. Cohen, R. W. Castenholz, and H. O. Halvorson), pp. 191–214. Alan R. Liss, Inc.
- Ward D. M., Brassell S. C., and Eglinton G. (1985) Archaeobacterial lipids in hot-spring microbial mats. *Nature* **318**, 656–659.
- Ward D. M., Shiea J., Zheng Y. B., Dobson G., Brassell S. C., and Eglinton G. (1989) Lipid biochemical markers and the composition of microbial mats. In *Microbial Mats: Physiological Ecology of Benthic Microbial Communities* (eds. Y. Cohen and E. Rosenberg), pp. 439–454. Am. Soc. Microbiol.
- Ward D. M., Santegoeds C. M., Nold S. C., Ramsing N. B., Ferris M. J., and Bateson M. M. (1997) Biodiversity within hot spring microbial mat communities: Molecular monitoring of enrichment cultures. *Antonie Van Leeuwenhoek Int. J. Gen. Molecul. Microbiol.* **71**, 143–150.
- Ward D. M., Ferris M. J., Nold S. C., and Bateson M. M. (1998) A

- natural view of microbial biodiversity within hot spring cyanobacterial mat communities. *Microbiol. Molecul. Biol. Rev.* **62**, 1353–1370.
- White D. C. (1988) Validation of quantitative analysis for microbial biomass, community structure, and metabolic activity. *Adv. Limnol.* **31**, 1–18.
- White D. C., Davis W. M., Nickels J. S., King J. D., and Bobbie R. J. (1979) Determination of the sedimentary microbial biomass by extractable lipid phosphate. *Oecologia* **40**, 51–62.
- Woese C. R. (1987) Bacterial evolution. *Microbiol. Rev.* **51**, 221–271.
- Yao M., Walker H. W., and Lillard D. A. (1970) Fatty acids from vegetative cells and spores of *Bacillus stearothermophilus*. *J. Bacteriol.* **102**, 877–878.
- Zeng Y. B., Ward D. M., Brassell S. C., and Eglinton G. (1992a) Biogeochemistry of hot spring environments. 2. Lipid compositions of Yellowstone (Wyoming, USA) cyanobacterial and *Chloroflexus* mats. *Chem. Geol.* **3-4**, 327–345.
- Zeng Y. B., Ward D. M., Brassell S. C., and Eglinton G. (1992b) Biogeochemistry of hot spring environments. 3. Apolar and polar lipids in the biologically active layers of a cyanobacterial mat. *Chem. Geol.* **3-4**, 347–360.
- Zhang C. L., Ye Q., Reysenbach A.-L., Goetz D., Peacock A., White D. C., Horita J., Cole D. R., Fang J., Pratt L., Fang J., and Huang Y. (2002) Carbon isotopic fractionations associated with thermophilic bacteria *Thermotoga maritima* and *Persephonella marina*. *Environ. Microbiol.* **4**, 58–64.
- Zhang C. L., Li Y., Ye Q., Fong J., Peacock A. D., Blunt E., Fang J., Lovley D. R., and White D. C. (2003) Carbon isotope signatures of fatty acids in *Geobacter metallireducens* and *Shewanella Algae*. *Chem. Geol.* **195**, 17–28.